

NEW DESCRIPTION OF SELF-DUAL METRICS

J. Tafel

Institute of Theoretical Physics, University of Warsaw, Hoża 69, 00-681 Warsaw, Poland,
email: tafel@fuw.edu.pl

Abstract. We show that Plebański's equation for self-dual metrics is equivalent to a pair of equations describing canonical transformations in 2-dimensional phase spaces. Examples of linearizations of these equations are given.

PACS number: 04.20(Cv, Jb) 02.40(Ky, Tt)

1 Introduction

Plebański [1] showed that complex metrics with the self-dual Riemann tensor [2], to be referred to as self-dual metrics, can be described in terms of one function Ω satisfying so-called Plebański's first equation

$$\Omega_{,q\tilde{q}}\Omega_{,p\tilde{p}} - \Omega_{,q\tilde{p}}\Omega_{,p\tilde{q}} = 1. \quad (1)$$

Given Ω the corresponding metric reads

$$g = \Omega_{,q\tilde{q}}dq d\tilde{q} + \Omega_{,p\tilde{p}}dp d\tilde{p} + \Omega_{,q\tilde{p}}dq d\tilde{p} + \Omega_{,p\tilde{q}}dp d\tilde{q}. \quad (2)$$

He also proposed an equivalent equation known as Plebański's second equation. At present there are several other descriptions of self-dual metrics [3, 4, 5, 6] (note that an erroronous sign in Husain's equation was corrected and an equivalence of the latter equation to (1) was proved [7]). Much attention was payed to possible complete integrability of the self-duality conditions (see [8] and references therein) .

In this paper we represent self-dual metrics in the form

$$g = \{u, v\}_{q\tilde{q}}dq d\tilde{q} + \{u, v\}_{p\tilde{p}}dp d\tilde{p} + \{u, v\}_{q\tilde{p}}dq d\tilde{p} + \{u, v\}_{p\tilde{q}}dp d\tilde{q}, \quad (3)$$

where the Poisson brackets are taken with respect to indicated coordinates, e.g. $\{u, v\}_{q\tilde{q}} = u_{,q}v_{,\tilde{q}} - u_{,\tilde{q}}v_{,q}$. Functions u and v are subject to the equations

$$\{u, v\}_{qp} = 1 \quad (4)$$

$$\{u, v\}_{\tilde{q}\tilde{p}} = 1. \quad (5)$$

Thus, u and v are related to q, p and to $\tilde{q}, \tilde{p}$ by canonical transformations. Solving (4) and (5) requires correlating these transformations. In section 3 we show that these equations can be linearized under some assumptions on functions generating the canonical transformations.

It is convenient to denote coordinates q, p by z^A , $A = 1, 2$, and coordinates $\tilde{q}, \tilde{p}$ by $\tilde{z}^A$. Partial derivatives of any function f with respect to these coordinates will be denoted by $f_{,A}$ and $f_{,\tilde{A}}$, respectively. Instead of $\{u, v\}_{z^A \tilde{z}^B}$ we will write $\{u, v\}_{A\tilde{B}}$. In this notation metric (2) reads

$$g = \Omega_{,A\tilde{B}} dz^A d\tilde{z}^B \quad (6)$$

and metric (3) reads

$$g = \{u, v\}_{A\tilde{B}} dz^A d\tilde{z}^B. \quad (7)$$

2 Equivalence of formulations

Let u and v satisfy equations (4) and (5). Then

$$\{u, v\}_{A[\tilde{B}, \tilde{C}]} = \{u, v\}_{\tilde{B}[A, C]} = 0, \quad (8)$$

hence there are functions Ω_A such that

$$\{u, v\}_{A\tilde{B}} = \Omega_{A, \tilde{B}} \quad (9)$$

and $\Omega_{[A, C]}$ do not depend on coordinates $\tilde{z}^A$. One can shift Ω_A by functions of z^A to achieve $\Omega_{[A, C]} = 0$. From this property it follows that there is a function Ω such that $\Omega_A = \Omega_{,A}$. Thus,

$$\{u, v\}_{A\tilde{B}} = \Omega_{,A\tilde{B}} \quad (10)$$

and metric (7) takes the form (6). Substituting (4), (5) and (10) into the identity

$$\{u, v\}_{q\tilde{q}} \{u, v\}_{p\tilde{p}} - \{u, v\}_{q\tilde{p}} \{u, v\}_{p\tilde{q}} = \{u, v\}_{qp} \{u, v\}_{\tilde{q}\tilde{p}} \quad (11)$$

shows that Ω satisfies Plebański's equation (1).

To prove that the reverse transformation exists is less straightforward. Let us use the antisymmetric tensors ϵ^{AB} and $\epsilon^{\tilde{A}\tilde{B}}$ to raise indices in a standard way, e.g. $p^A = \epsilon^{AB} p_{,B}$. This notation allows to write equations (4) and (5) in the form

$$v^A u_{,A} = 1 \quad (12)$$

$$v^{\tilde{A}} u_{,\tilde{A}} = 1. \quad (13)$$

Assume that a function Ω satisfies the equation

$$\Omega_{,A\tilde{B}} \Omega_{,C}^{\tilde{B}} = \epsilon_{AC}, \quad (14)$$

which is equivalent to (1). Our aim is to prove the existence of solutions u, v of equations (10), (12) and (13).

Taking contractions of (10) with v^A or u^A and using (12) we can replace equations (10) by

$$v_{,\tilde{B}} = v^A \Omega_{,A\tilde{B}} \quad (15)$$

and

$$u_{,\tilde{B}} = u^{,A} \Omega_{,A\tilde{B}}. \quad (16)$$

It follows from (14)-(16) that

$$v^{,\tilde{B}} u_{,\tilde{B}} = v^{,A} u^{,C} (\Omega_{,A}^{,\tilde{B}} \Omega_{,C\tilde{B}}) = v^A u_A. \quad (17)$$

Thus, equation (13) is satisfied if equations (12) and (14)-(16) are satisfied.

Let us write equations (15) and (16) as

$$D_{\tilde{B}} v = 0 \quad (18)$$

$$D_{\tilde{B}} u = 0, \quad (19)$$

where operators $D_{\tilde{B}}$ are given by

$$D_{\tilde{B}} = \partial_{\tilde{B}} + \Omega^{,A}_{\tilde{B}} \partial_A. \quad (20)$$

Vectors $D_{\tilde{B}}$ are part of a null basis for metric (2). In view of twistor constructions related to self-dual metrics [9] it is not suprising that $D_{\tilde{B}}$ commute provided equation (14) is satisfied,

$$[\partial_{\tilde{B}} + \Omega^{,A}_{\tilde{B}} \partial_A, \partial_{\tilde{C}} + \Omega^{,D}_{\tilde{C}} \partial_D] = 2\Omega^{,A}_{[\tilde{B}} \Omega^{,D}_{\tilde{C}]} \partial_D = 2(\Omega^{,A}_{[\tilde{B}} \Omega^{,D}_{\tilde{C}]})_{,A} \partial_D = 0. \quad (21)$$

Let v be a solution of (18). It remains to solve equations (12) and (19), which form a linear system for a function u . The system is integrable since all the operators $D_{\tilde{B}}$, $v^{,A} \partial_A$ commute due to (21) and the equality

$$[\partial_{\tilde{B}} + \Omega^{,C}_{\tilde{B}} \partial_C, v^{,A} \partial_A] = (v_{,\tilde{B}} + \Omega^{,C}_{\tilde{B}} v_{,C})^{,A} \partial_A = 0 \quad (22)$$

forced by (18). Thus, equations (10), (12), (13) have solutions and we obtain the following theorem.

Theorem 1. *All complex self-dual metrics can be locally represented in the form (3), where u and v are subject to equations (4) and (5).*

Note that if u , v and all the coordinates are real, then metric (3) is real with the signature $++--$. The same signature is obtained when u and iv are real and

$$\tilde{q} = \epsilon \bar{q}, \quad \tilde{p} = -\epsilon \bar{p}, \quad \epsilon = \pm 1, \quad (23)$$

where the bar denotes the complex conjugate. Simple conditions of this kind are not available for the Euclidean signature of g . This difficulty appears also in other approaches to self-duality [5, 6].

Equations (4) and (5) possess point symmetries which are easy to obtain if we denote u , v by f^A and write equations (4), (5) as

$$\det\left(\frac{\partial f^A}{\partial z^B}\right) = \det\left(\frac{\partial f^A}{\partial \bar{z}^B}\right) = 1. \quad (24)$$

Let new variables f'^A be functions of f^B , new coordinates z'^A be functions of z^B and $\tilde{z}'^A$ be functions of $\tilde{z}^B$. This transformation preserves equations (24) provided

$$\det\left(\frac{\partial f'^A}{\partial f^B}\right) = \det\left(\frac{\partial z'^A}{\partial z^B}\right) = \det\left(\frac{\partial \tilde{z}'^A}{\partial \tilde{z}^B}\right) = c, \quad (25)$$

where c is a constant. In terms of Poisson brackets equations (25) read

$$\{u', v'\}_{uv} = \{q', p'\}_{qp} = \{\tilde{q}', \tilde{p}'\}_{\tilde{q}\tilde{p}} = c. \quad (26)$$

There are also natural discrete symmetries given by an appropriate interchange of variables. All above transformations preserve the metric modulo a constant factor.

3 Reductions to linear equations

In terms of a generating function $S(v, q, \tilde{q}, \tilde{p})$, where $\tilde{q}, \tilde{p}$ appear as parameters, solutions u, v of (4) are given implicitly by

$$p = S_{,q}, \quad u = S_{,v}. \quad (27)$$

Similarly, equation (5) can be formally solved by means of a generating function $\tilde{S}(v, \tilde{q}, q, p)$,

$$\tilde{p} = \tilde{S}_{,\tilde{q}}, \quad u = \tilde{S}_{,v}. \quad (28)$$

Functions S and $\tilde{S}$ should be correlated in such a way that solutions u, v obtained from (27) or (28) coincide.

Assume that S and $\tilde{S}$ are linear in $\tilde{p}$ and p , respectively,

$$S = S'(v, q, \tilde{q}) + \tilde{p}a(v, q, \tilde{q}), \quad \tilde{S} = \tilde{S}'(v, q, \tilde{q}) + p\tilde{a}(v, q, \tilde{q}). \quad (29)$$

Substituting (29) into (27) and (28) leads to the following conditions guaranteeing the uniqueness of u and v

$$\tilde{a}_{,\tilde{q}} = a_{,q}^{-1}, \quad \tilde{a}_{,v} = a_{,v}a_{,q}^{-1} \quad (30)$$

$$S'_{,q} + a_{,q}\tilde{S}'_{,\tilde{q}} = 0, \quad S'_{,v} - \tilde{S}'_{,v} + a_{,v}\tilde{S}'_{,\tilde{q}} = 0. \quad (31)$$

Equations (30) form a nonlinear system for the functions a and $\tilde{a}$. Given a and $\tilde{a}$ it remains to solve linear equations (31) for functions S' and $\tilde{S}'$.

Equations (30) are integrable with respect to $\tilde{a}$ provided a satisfies the following quasilinear equation

$$a_{,vq} + a_{,q}a_{,v\tilde{q}} - a_{,v}a_{,q\tilde{q}} = 0. \quad (32)$$

Simple solutions of (32) can be found by assuming that one of the second derivatives of a vanishes. For instance, for $a_{,vq} = 0$ one obtains $a = \tilde{q}(v + q)$. For this function a equations (31) yield

$$S = F_{,q} + \tilde{p}\tilde{q}(v + q), \quad (33)$$

where function $F(v, q, \tilde{q})$ is subject to the linear equation

$$F_{,vq} + F_{,v\tilde{q}} - F_{,q\tilde{q}} = 0. \quad (34)$$

In this case the corresponding metric (3) possesses the translational Killing vector $\partial_p - \partial_{\tilde{p}}$. For this reason it must belong to the class of generalized Hawking metrics [6].

If $a_{,v\tilde{q}} = 0$ then the generating function S can be put into the form

$$S = F_{,v} - e^q F_{,\tilde{q}} + \tilde{p}(q\tilde{q} + e^q v + c(q)). \quad (35)$$

Here c is an arbitrary function of q , with its derivative denoted by $\dot{c}$, and F has to satisfy the linear equation

$$F_{,vq} + (\tilde{q} + e^q v + \dot{c})F_{,v\tilde{q}} - e^q F_{,q\tilde{q}} - e^q F_{,\tilde{q}} = 0. \quad (36)$$

If $a_{,q\tilde{q}} = 0$ one obtains

$$S = F_{,\tilde{q}} + \tilde{p}(v\tilde{q} + e^{-v} q + c(v)) \quad (37)$$

and the following linear equation for F

$$e^{-v} F_{,v\tilde{q}} + (e^{-v} q - \tilde{q} - \dot{c})F_{,q\tilde{q}} + F_{,vq} + F_{,q} = 0. \quad (38)$$

Given a solution F of equation (36) or (38) one can find functions u, v from relations (27) and then construct the corresponding self-dual metric (3). Up to the best knowledge of the author the reduction of the Plebański equation, for some ansatz, to the linear equations (36) or (38) is new.

4 Conclusions

We have shown that all self-dual metrics can be represented in the form (3), where functions u, v have to satisfy equations (4) and (5) defining canonical transformations in a 2-dimensional phase space.

The new formulation creates new possibilities of construction of self-dual metrics. Equations (4), (5) are equivalent to linear ones (34), (36) or (38) under the assumption that functions u, v are given by (27) and S takes the form (33), (35) or (37). An open problem is whether these linear reductions admit interesting solutions other than gravitational instantons of Hawking [10].

In our opinion the new formulation can be also useful as a new tool in classification of completely integrable equations. It is well known that many of these equations can be considered as reductions of the self-dual Yang-Mills equations [11] or the self-dual Einstein equations [12] (see also [8]). Perhaps, due to a new form of self-duality conditions, one can obtain in this way further completely integrable equations.

Acknowledgements. This work was partially supported by the Polish Committee for Scientific Research (grant 1 P03B 075 29).

References

- [1] Plebański J 1975 *J. Math. Phys.* **16** 2395
- [2] Penrose R 1976 *Gen. Rel. Grav.* **7** 31
- [3] Ashtekar A, Jacobson T and Smolin L 1988 *Commun. Math. Phys.* **115** 631
- [4] Chakravarty S, Mason L J and Newman E T 1991 *J. Math. Phys.* **32** 1458
- [5] Grant J D E 1993 *Phys. Rev.* **D48** 2606
- [6] Husain V 1994 *Class. Quantum Grav.* **11** 927
- [7] Jakimowicz M and Tafel J 2005 *work in preparation*
- [8] Mason L J and Woodhouse N M J 1996 *Integrability, self-duality and twistor theory* (Clarendon press, Oxford)
- [9] Penrose R and Rindler W 1986 *Spinors and spacetime*, vol. II (Cambridge University Press, Cambridge)
- [10] Hawking S W 1977 *Phys. Let.* **60A** 81
- [11] Ward R S 1985 *Phil. Trans. R. Soc.* **A315** 451
- [12] Dunajski M, Mason L J and Woodhouse N M J 1998 *J. Phys. A: Math. Gen.* **31** 6019